

Detection of mitochondrial ROS levels using flow cytometry

XZ Xiaowei Zheng SC Sergiu-Bogdan Catrina

Updated date: Apr 3, 2023

 An abbreviated version of this protocol was published in eLIFE in Feb 2022

Repression of hypoxia-inducible factor-1 contributes to increased mitochondrial reactive oxygen species production in diabetes

DOI: 10.7554/eLife.70714

Related files

 Detection of mitochondrial ROS levels using Flow Cytometry.docx



How to cite: (Readers should cite both the Bio-protocol preprint and the original research article where this protocol was used)

1. Zheng, X. and Catrina, S. (2023). Detection of mitochondrial ROS levels using flow cytometry. Bio-protocol Preprint. bio-protocol.org/prep2187.
2. Zheng, X., Narayanan, S., Xu, C., Eliasson Angelstig, S., Grünler, J., Zhao, A., Di Toro, A., Bernardi, L., Mazzone, M., Carmeliet, P., Del Sole, M., Solaini, G., Forsberg, E. A., Zhang, A., Brismar, K., Schiffer, T. A., Rajamand Ekberg, N., Botusan, I. R., Palm, F. and Catrina, S. (2022). Repression of hypoxia-inducible factor-1 contributes to increased mitochondrial reactive oxygen species production in diabetes. eLIFE. DOI: [10.7554/eLife.70714](https://doi.org/10.7554/eLife.70714)

Copyright: Content may be subjected to copyright.